## **What You Should Know**

#### Will this medicine work for me?

- The antidepressants presented in this decision aid all work the same for treating depression.
- Most people with depression can find one that can make them feel better.
- 6 out of 10 people will feel better with the first antidepressant they try and the rest will have to try other antidepressants before they find the one that is right for them.

### How long before I feel better?


 Most people need to take an antidepressant regularly for at least 6 weeks to begin to get the full effect.

### **Understanding side effects**

- Most people taking antidepressants have at least one side effect.
- Many side effects go away after a few weeks,
   but some only go away after you stop the medicine.


# **Weight Change**

Some people may experience weight change. It is most likely to occur over six to twelve months and depends on your actual weight. The chart below is based on a 150 lb person.




## **Sexual Issues**

Some people may experience loss of sexual desire (libido) or loss of ability to reach orgasm because of their antidepressant.




# Sleep

Some people may experience sleepiness or insomnia because of their antidepressant.




## Cost

These figures are estimates and are for comparative reference only. Actual out-of-pocket costs vary over time, by pharmacy, insurance plan coverage, preparation and dosage.



## **Stopping Approach**

Quitting your medicine all at once can make you feel sick, as if you had the flu (e.g. headache, dizziness, light-headedness, nausea or anxiety).



## **Keep in Mind**

### Depression medicines may cause some:

- · constipation, diarrhea and nausea
- increased risk of suicidal thoughts and behaviors (18- to 24-year-olds)
- · harm to an unborn child
- · risk of developing serotonin syndrome, a potentially life-threatening condition
- possible drug-drug interactions

#### **Additional considerations**

## Citalopram (Celexa®)

Can cause problems with your heart

**Escitalopram** 

Currently no other issues

Fluoxetine

(Lexapro®)

More likely to interact with other drugs you are taking

**Fluvoxamine** 

More likely to cause constipation, diarrhea or nausea

(Luvox®)

(Prozac®)

Not officially recognized as a treatment for Major Depressive Disorder

**Paroxetine** 

If you are pregnant, this medicine is more likely to cause

problems with your unborn child

Sertraline

More likely to cause diarrhea

(Zoloft®)

(Paxil®)

**Desvenlafaxine** 

Tell your doctor if you have high blood pressure

(Pristiq®)

**Duloxetine** Can help with pain

(Cymbalta®) Tell your doctor if you have high blood pressure

Venlafaxine

More likely to cause nausea and vomiting

(Effexor®) Can cause problems with your heart

Tell your doctor if you have high blood pressure

Bupropion (Wellbutrin®)

Higher risk of seizures

Mirtazapine

Starts to work more quickly

(Remeron®)

Amiptriptyline or Nortriptyline

More likely to cause constipation, diarrhea or nausea

Can help with pain

(Elavil® or Aventyl HCl®)

If you are elderly, this medication may not be the best option

SSRI

SNRIS

**TCAs**